

On the possibility of a measurement of the CP Majorana phase in the $0\nu\beta\beta$ -decay

F. Šimkovic,^{1,2} S.M. Bilenky,^{2,3} Amand Faessler,⁴ and Th. Gutsche⁴

¹*Department of Nuclear Physics and Biophysics, Comenius University,
Mlynská dolina F1, SK-842 15 Bratislava, Slovakia*

²*Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Moscow region, Russia*

³*TRIUMF 4004 Wesbrook Mall, Vancouver BC, V6T 2A3 Canada*

⁴*Institute of Theoretical Physics, University of Tuebingen, 72076 Tuebingen, Germany*

In view of recent measurements of the mixing angle θ_{13} a possibility to determine the difference of two CP Majorana phases of the neutrino mixing matrix from the study of neutrinoless double-beta decay is investigated. We show that in the case of the inverted hierarchy of neutrino masses it might be possible if neutrinoless double-beta decay will be observed. The required experimental accuracies and uncertainty in the calculated nuclear matrix elements of the process are discussed.

PACS numbers: 23.40.Hc, 21.60.Jz, 14.60.St, 12.60.Jv

I. INTRODUCTION

The discovery of neutrino oscillations in the SuperKamiokande (atmospheric neutrinos) [1], SNO (solar neutrinos) [2], KamLAND (reactor neutrinos) [3], MINOS [4] (accelerator neutrinos) and other neutrino experiments gives us compelling evidence that neutrinos possess small masses and flavor neutrino fields are mixed.

Neutrino flavor states $|\nu_\alpha\rangle$ ($\alpha = e, \mu, \tau$) are connected to the states of neutrinos with masses m_j ($|\nu_j\rangle$) by the following standard mixing relation

$$|\nu_\alpha\rangle = \sum_{j=1}^3 U_{\alpha j}^* |\nu_j\rangle \quad (\alpha = e, \mu, \tau), \quad (1)$$

where U is the 3×3 Pontecorvo-Maki-Nakagawa-Sakata (PMNS) unitary mixing matrix [5]. If massive neutrinos are Dirac (Majorana) particles in this case the PMNS matrix contains one (three) CP phases.

The problem of the determination of the CP phases is one of the major challenges in today's neutrino physics. Some information about lepton phases could help to solve the problem of the baryon asymmetry of the Universe [6]. The discovery in the Daya Bay [7], RENO [8], T2K [9] and Double Chooz [10] experiments that the angle θ_{13} is relatively large opened a possibility of the measurement of the Dirac phase δ in long baseline accelerator experiments. The Majorana CP phases can be determined only through the observation of neutrinoless double β -decay ($0\nu\beta\beta$ -decay) [11–13].

The CP phases enter into the effective Majorana mass [14, 15, 17] defined as

$$m_{\beta\beta} = \sum_j U_{ej}^2 m_j. \quad (2)$$

This quantity depends also on the neutrino oscillation parameters $\theta_{12}, \theta_{13}, \Delta m_{\text{SUN}}^2, \Delta m_{\text{ATM}}^2$, the lightest neutrino mass and the type of the neutrino mass spectrum (normal or inverted).

The effective Majorana mass $m_{\beta\beta}$ can be determined in the neutrinoless double beta decay of even-even nuclei [18, 19]

$$(A, Z) \rightarrow (A, Z + 2) + 2e^- \quad (3)$$

by converting the $0\nu\beta\beta$ -decay half-life to $|m_{\beta\beta}|$ using calculated nuclear matrix elements (NMEs).

The goal of this paper is to discuss a possibility for determining the Majorana CP phases from data of $0\nu\beta\beta$ experiments of the next generation assuming that the $0\nu\beta\beta$ -decay will be observed in these experiments. The problem of the experimental accuracies and uncertainties of the calculated NMEs will also be addressed.

II. NEUTRINO OSCILLATIONS AND EFFECTIVE MAJORANA MASS

It was proved in the experiments with atmospheric, solar, reactor and accelerator neutrinos that flavor neutrinos oscillate from one flavor (electron-, muon-, and tau-) to another due to neutrino mixing and non-zero neutrino mass-squared differences. All existing neutrino oscillation data (with the exception of the LSND [20], MiniBooNE [21], short baseline reactor [22] and Gallium [23] anomalies) are perfectly described by the minimal scheme of three-neutrino mixing.

In the case of Dirac neutrinos the unitary 3×3 PMNS neutrino mixing matrix can be parameterized as follows

$$U = \begin{pmatrix} c_{12}c_{13} & c_{13}s_{12} & e^{-i\delta}s_{13} \\ -c_{23}s_{12} - e^{i\delta}c_{12}s_{13}s_{23} & c_{12}c_{23} - e^{i\delta}s_{12}s_{13}s_{23} & c_{13}s_{23} \\ s_{12}s_{23} - e^{i\delta}c_{12}c_{23}s_{13} & -e^{i\delta}c_{23}s_{12}s_{13} - c_{12}s_{23} & c_{13}c_{23} \end{pmatrix} \quad (4)$$

where $c_{ij} \equiv \cos \theta_{ij}$, $s_{ij} \equiv \sin \theta_{ij}$. θ_{12}, θ_{13} and θ_{23} are mixing angles and δ is the CP phase. If neutrinos are Majorana particles the matrix U in Eq. (4) is multiplied by a diagonal phase matrix $P = \text{diag}(e^{i(\alpha_1/2-\delta)}, e^{i(\alpha_2/2-\delta)}, 1)$, which contains two additional CP phases α_1 and α_2 .

With the discovery of neutrino oscillations we know:

- The values of the large mixing angles θ_{12} and θ_{23} . The value of the relatively small angle θ_{13} measured recently in the Double Chooz [10], the Daya Bay [7] and RENO [8] reactor neutrino experiments.




FIG. 1. (Color online) The CP phase α_2 (left panels) and difference of phases α_{21} (right panels) are plotted, respectively, as function of the absolute value of the effective majorana mass for the IH and NH of neutrino masses and chosen value of m_0 .

- The solar and atmospheric mass-squared differences¹ $\Delta m_{\text{SUN}}^2 = \Delta m_{12}^2$ and $\Delta m_{\text{ATM}}^2 = \Delta m_{23}^2$ (Normal spectrum), $\Delta m_{\text{ATM}}^2 = -\Delta m_{13}^2$ (Inverted spectrum).

We do not know the value of the lightest neutrino mass, the CP phases and the character of the neutrino mass spectrum (normal or inverted).

From the data of the MINOS experiment [4] it was found that $\Delta m_{\text{ATM}}^2 = (2.43 \pm 0.13) \times 10^{-3} \text{ eV}^2$. From the analysis of the KamLAND and solar data it was obtained that $\tan^2 \theta_{12} = 0.452_{-0.033}^{+0.035}$ [3]. From the global fit to all data it was inferred that [24] $\Delta m_{\text{SUN}}^2 = (7.65_{-0.20}^{+0.13}) \times 10^{-5} \text{ eV}^2$ and $\sin^2 \theta_{23} = 0.50_{-0.06}^{+0.07}$. Finally from the analysis of the Daya Bay[7] and RENO data[8] one obtains $\sin^2 2\theta_{13} = 0.092 \pm 0.016 \text{ (stat)} \pm 0.005 \text{ (syst)}$ and $\sin^2 2\theta_{13} = 0.103 \pm 0.013 \text{ (stat)} \pm 0.011 \text{ (syst)}$, respectively.

The effective Majorana mass is given by

$$|m_{\beta\beta}| = |c_{12}^2 c_{13}^2 e^{i\alpha_1} m_1 + s_{12}^2 c_{13}^2 e^{i\alpha_2} m_2 + s_{13}^2 m_3| \quad (5)$$

or by the full expression

$$\begin{aligned} |m_{\beta\beta}|^2 = & c_{12}^4 c_{13}^4 m_1^2 + s_{12}^4 c_{13}^4 m_2^2 + s_{13}^4 m_3^2 \\ & + 2c_{12}^2 s_{12}^2 c_{13}^4 m_1 m_2 \cos(\alpha_1 - \alpha_2) \\ & + 2c_{12}^2 c_{13}^2 s_{13}^2 m_1 m_3 \cos \alpha_1 + 2s_{12}^2 c_{13}^2 s_{13}^2 m_2 m_3 \cos \alpha_2. \end{aligned} \quad (6)$$

From this equation it simply follows that the effective Majorana mass depends on the character of the neutrino mass spectrum and three unknown parameters: the lightest neutrino mass and two CP phases.

In the three-neutrino case two mass spectra are currently possible:

- Normal Spectrum (NS): $m_1 < m_2 < m_3$: $\Delta m_{12}^2 \ll \Delta m_{23}^2$. In this case

$$m_2 = \sqrt{\Delta m_{\text{SUN}}^2 + m_0^2}, \quad m_3 = \sqrt{\Delta m_{\text{ATM}}^2 + \Delta m_{\text{SUN}}^2 + m_0^2}$$

with $m_0 = m_1$.

¹ We use the following definition $\Delta m_{ij}^2 = m_j^2 - m_i^2$

- Inverted Spectrum (IS), $m_3 < m_1 < m_2$: $\Delta m_{12}^2 \ll |\Delta m_{13}^2|$. We have

$$m_1 = \sqrt{\Delta m_{\text{ATM}}^2 + m_0^2}, \quad m_2 = \sqrt{\Delta m_{\text{ATM}}^2 + \Delta m_{\text{SUN}}^2 + m_0^2}$$

with $m_0 = m_3$

Here, $m_0 = m_1(m_3)$ is the lightest neutrino mass.

We will consider the following two neutrino mass hierarchies:

1. Normal Hierarchy (NH): $m_1 \ll m_2 \ll m_3$:

In this case for the neutrino masses we have

$$m_1 \ll \sqrt{\Delta m_{\text{SUN}}^2}, \quad m_2 \simeq \sqrt{\Delta m_{\text{SUN}}^2}, \quad m_3 \simeq \sqrt{\Delta m_{\text{ATM}}^2}.$$

Neglecting the negligibly small contribution of m_1 we find

$$\cos \alpha_2 \simeq \frac{|m_{\beta\beta}|^2 - s_{12}^4 c_{13}^4 \Delta m_{\text{SUN}}^2 - s_{13}^4 \Delta m_{\text{ATM}}^2}{2s_{12}^2 c_{13}^2 s_{13}^2 \sqrt{\Delta m_{\text{SUN}}^2 \Delta m_{\text{ATM}}^2}}. \quad (7)$$

For the effective Majorana mass we then have the following range of values

$$|s_{12}^2 c_{13}^2 \sqrt{\Delta m_{\text{SUN}}^2} - s_{13}^2 \sqrt{\Delta m_{\text{ATM}}^2}| \leq |m_{\beta\beta}| \leq s_{12}^2 c_{13}^2 \sqrt{\Delta m_{\text{SUN}}^2} + s_{13}^2 \sqrt{\Delta m_{\text{ATM}}^2}. \quad (8)$$

Using the best-fit values of the mass squared differences and the mixing angles we find

$$1.5 \text{ meV} \leq |m_{\beta\beta}| \leq 3.8 \text{ meV}$$

2. Inverted Hierarchy (IH): $m_3 \ll m_1 < m_2$:

In the IH scenario $m_3 \ll \sqrt{\Delta m_{\text{ATM}}^2}$ and $m_1 \simeq m_2 \simeq \sqrt{\Delta m_{\text{ATM}}^2}$. We find

$$\cos \alpha_{12} = \frac{|m_{\beta\beta}|^2 - c_{13}^4 (1 - 2s_{12}^2 c_{12}^2) \Delta m_{\text{ATM}}^2}{2c_{12}^2 s_{12}^2 c_{13}^4 \Delta m_{\text{ATM}}^2} \quad (9)$$

where $\alpha_{12} = \alpha_1 - \alpha_2$. For the absolute value of the effective Majorana mass we find the range

$$|\cos 2\theta_{12}| c_{13}^2 \sqrt{\Delta m_{\text{ATM}}^2} \leq |m_{\beta\beta}| \leq c_{13}^2 \sqrt{\Delta m_{\text{ATM}}^2}. \quad (10)$$

Using the best-fit values of the parameters we find the following range for $|m_{\beta\beta}|$ in the case of the IH:

$$18 \text{ meV} \leq |m_{\beta\beta}| \leq 48 \text{ meV}.$$

The absolute value of the neutrino mass can be determined from the precise measurement of the end-point part of the β -spectrum of the tritium [25] and other β -decay measurements [26]. Cosmological observations allow to infer the sum of the neutrino masses

$$m_{\text{cosmo}} = \sum_k^3 m_k. \quad (11)$$

The current limits on m_{cosmo} depend on the type of observations included in the fit [27]. The CMB primordial gives $\leq 1.3 \text{ eV}$, CMB+distance $\leq 0.58 \text{ eV}$, galaxy distribution and lensing of galaxies $\leq 0.6 \text{ eV}$. On the other hand the largest photometric red shift survey yields $\leq 0.28 \text{ eV}$ [28]. It is expected that future cosmological observables will be sensitive to m_{cosmo} in the range (0.05-0.1) eV (see, e.g., the recent summary [27]). Thus, the inverted neutrino mass hierarchy will be tested by future precision cosmology. In the case of the inverted mass hierarchy we have for the sum of the neutrino masses

$$m_{\text{cosmo}} \simeq 2\sqrt{\Delta m_{\text{ATM}}^2} \simeq 100 \text{ meV}. \quad (12)$$

So, there is a chance that future cosmological observation apparently could restrict the value of m_0 below 10 meV.

In Fig. 1, by exploiting Eq. (6), the Majorana CP phase α_2 (or difference of phases $\alpha_{21} = \alpha_2 - \alpha_1$) is plotted as function of the absolute value of the effective Majorana mass for chosen values of m_0 and by assuming the NH (IH) of




FIG. 2. (Color online) The difference of CP phases $\alpha_{21} = \alpha_2 - \alpha_1$ plotted as function of $|m_{\beta\beta}|^{obs}$ for the inverted hierarchy of neutrino masses. The current experimental errors of neutrino mixing parameters and mass squared differences are taken into account; m_0 is taken to be in the range of 0 to 10 meV. The blue, red and orange regions correspond to $|m_{\beta\beta}|^{obs}$ with error $(\sigma_{\beta\beta}/|m_{\beta\beta}|^{obs})$ of 0%, 15% and 25%, respectively.

neutrino masses. The second phase α_1 is considered to be arbitrary. We found that for m_0 within a range of 0 to 10 meV α_{21} depends only weakly on $|m_{\beta\beta}|$, if the IH is considered. It is due to the fact that the third term on the right hand side of Eq. (5) is small in comparison with the first two terms. A different situation occurs in the case of the NH. Results depend strongly on m_0 , where its value cannot possibly be determined from other experiments or cosmology, if it is lower than a few meV. Thus, by measuring $0\nu\beta\beta$ -decay it will not be possible to obtain a neutrino mass model independent information on the value of at least one of the three CP Majorana phases, if there is a normal hierarchy of neutrino masses. In some particle physics models there is a possibility to predict all three neutrinos masses. In these cases values for one of the CP Majorana phases can be obtained when considering the second phase to be arbitrary and by observing the $0\nu\beta\beta$ -decay. However, the results depend strongly on the value of m_0 and the type of the neutrino mass hierarchy, normal or inverted. It is interesting to note that for $m_0 = 3$ meV the minimal value of $|m_{\beta\beta}|$ does not appear for the case of CP-conservation (see left upper panel of Fig. 1) but for $\alpha_2 \simeq 0.79\pi$.

III. EFFECTIVE MAJORANA MASS AND THE $0\nu\beta\beta$ -DECAY

Assuming that the $0\nu\beta\beta$ -decay is driven by the Majorana neutrino mass mechanism, we have for the effective Majorana mass $m_{\beta\beta}$

$$|m_{\beta\beta}| = \frac{1}{\sqrt{T_{1/2}^{0\nu} G_{0\nu}(Q_{\beta\beta}, Z) |M'^{0\nu}|}}. \quad (13)$$

Here, $T_{1/2}^{0\nu}$, $G_{0\nu}(Q_{\beta\beta}, Z)$ and $M'^{0\nu}$ are, respectively, half-life of the $0\nu\beta\beta$ -decay, the known phase-space factor and the nuclear matrix element, which depends on nuclear structure.

The future experiments, CUORE (^{130}Te), EXO, KamLAND-Zen (^{136}Xe), MAJORANA/GERDA (^{76}Ge), SuperNEMO (^{82}Se), SNO+ (^{150}Nd), and others [18], with a sensitivity

$$|m_{\beta\beta}| \simeq \text{a few } 10^{-2} \text{ eV} \quad (14)$$

will probe the IH of neutrino masses. In the case of the normal mass hierarchy $|m_{\beta\beta}|$ is much too small in order that $0\nu\beta\beta$ -decay will be detected in experiments of the next generation.

If the $0\nu\beta\beta$ -decay will be observed the measured half-life $T_{1/2}^{0\nu-exp}$ with experimental error σ_{exp} can be converted into an “observed effective Majorana mass” $|m_{\beta\beta}|^{obs}$ and its error $\sigma_{\beta\beta}$ as

$$\frac{\sigma_{\beta\beta}}{|m_{\beta\beta}|^{obs}} = \sqrt{\frac{1}{4} \left(\frac{\sigma_{exp}}{T_{1/2}^{0\nu-obs}} \right)^2 + \left(\frac{\sigma_{th}}{|M'^{0\nu}|} \right)^2}. \quad (15)$$

Here, σ_{th} is a “theoretical error” of the nuclear matrix element $|M'^{0\nu}|$.

In Fig. 2 we plot the difference of the CP Majorana phases α_{21} as function of $|m_{\beta\beta}|^{obs}$ for the inverted hierarchy of neutrino masses and by assuming 0% (blue region), 15% (red region) and 25% uncertainty (orange region) in $|m_{\beta\beta}|^{obs}$. The current experimental errors of neutrino mixing parameters and mass squared differences are taken into account. The lightest neutrino mass m_0 is assumed to be within a range from 0 to 10 meV and one of the CP violating phases α_1 (or α_2) is taken to be arbitrary. We see that if the considered accuracies are achieved it can be possible to determine the value of the CP phase difference α_{12} . However, for $\sigma_{\beta\beta}/|m_{\beta\beta}|^{obs} > 50\%$ it will be difficult, or impossible, to gain reliable information about the value of α_{12} .

The uncertainty σ_{th} in the calculated $0\nu\beta\beta$ decay NME is a complicated and more involved problem. $M'^{0\nu}$ consists of the Fermi (F), Gamow-Teller (GT) and tensor (T) parts [29–31]:

$$M'^{0\nu} = \left(\frac{g_A^{eff}}{g_A} \right)^2 \left(-\frac{M_F^{0\nu}}{(g_A^{eff})^2} + M_{GT}^{0\nu} - M_T^{0\nu} \right) \quad (16)$$

Here, $g_A^{eff}(g_A)$ is the quenched (unquenched) axial-vector coupling constant. $M'^{0\nu}$ is a function of $(g_A^{eff})^2$, which appears in the Fermi matrix element and also enters in the calculation of the Gamow-Teller and tensor constituents due to a consideration of the nucleon weak-magnetism terms [32]. This definition of $M'^{0\nu}$ [29–31] allows to display the effects of the uncertainties in g_A^{eff} and to use the same phase factor $G^{0\nu}$ when calculating the $0\nu\beta\beta$ -decay rate.

The treatment of quenching g_A^{eff} is an important source of difference between the calculated $0\nu\beta\beta$ -decay NMEs [30, 33]. Quenching of the axial-vector coupling was introduced to account for the fact that the calculated strengths of the Gamow-Teller β -decay transitions to individual final states are significantly larger than the experimental ones. Formally this is accomplished by replacing the true vacuum value of the coupling constant $g_A = 1.269$ by a quenched value $g_A^{eff} = 1.0$. It is not clear whether a similar phenomenon exists for other multipoles besides $J = 1^+$.

Different nuclear structure methods have been used for the calculation of $M'^{0\nu}$, in particular, interacting shell model (ISM) [34, 35], quasiparticle random phase approximation (QRPA) [29–31, 36], projected Hartree-Fock Bogoliubov approach (PHFB, PQQ2 parametrization) [37], energy density functional method (EDF) [38] and interacting boson model (IBM) [39]. By assuming an unquenched g_A the ISM values of the NMEs are about a factor 2-3 smaller than the NMEs of other approaches (see Table 3 of [19]). The results of the QRPA, IBM, EDF and PHB approaches differ by a factor less than two. Their results agree rather well with each other in the case of the $0\nu\beta\beta$ -decay of ^{130}Te .

A detailed study of uncertainties in the calculated $0\nu\beta\beta$ -decay NMEs was performed within the QRPA approach [29–31, 36]. The average matrix element $\langle M'^{0\nu} \rangle$ (averaged over different NN potentials, choices of the single particle space, variants of the QRPA approach) was evaluated, as well as its variance σ :

$$\sigma^2 = \frac{1}{N-1} \sum_{i=1}^N \left(M'_i{}^{0\nu} - \langle M'^{0\nu} \rangle \right)^2. \quad (17)$$

Further progress was achieved by performing a self-consistent calculation of the NMEs in which the pairing and residual interactions as well as the two-nucleon short-range correlations were derived from the same modern nucleon-nucleon potentials, namely, from the charge-dependent Bonn potential (CD-Bonn) and the Argonne V18 potential [31]. The particle-particle strength was adjusted to the $2\nu\beta\beta$ -decay half-life eliminating one of the main reasons for variability of the calculated $M'^{0\nu}$ within the QRPA-like methods [29, 36].

Here, we update the calculation of the average $\langle M'^{0\nu} \rangle$ and its variance σ for the $0\nu\beta\beta$ -decay of ^{76}Ge and ^{136}Xe . The recommended half-life value of $2\nu\beta\beta$ -decay of ^{76}Ge [40] and the recently measured half-life of the $2\nu\beta\beta$ -decay of ^{136}Xe [41, 42] were considered. The calculations were performed for the CD-Bonn and Argonne potentials, 3 different sizes of the model space, a quenched and unquenched axial vector coupling constant and two variants of the QRPA - the standard QRPA and the renormalized QRPA (RQRPA). Following values were found:

$$\begin{aligned} \langle M'^{0\nu} \rangle &= 4.78, & \sigma &= 0.74 & (^{76}\text{Ge}) \\ &= 2.33, & \sigma &= 0.36 & (^{136}\text{Xe}). \end{aligned} \quad (18)$$




FIG. 3. (Color online) The spread of the NMEs for the $0\nu\beta\beta$ -decay of ^{76}Ge and ^{136}Xe calculated within QRPA and RQRPA by assuming Argonne (A) and CD-Bonn (B) NN interactions, 3 different sizes of the model space and quenched ($g_A^{eff} = 1.00$) and unquenched ($g_A^{eff} = g_A = 1.269$) weak coupling constant g_A .

The nuclear radius was assumed to be $R=1.2 A^{1/3}$ fm. We see that the variance σ is about 15% of the full NME $M'^{0\nu}$. Of course, these results are valid only for the QRPA approach and the considered averaging scheme.

The calculated sets of 24 NMEs for ^{76}Ge and of 24 NMEs for ^{136}Xe are presented in Fig. 3. The results do not depend much on the size of the model space and on the type of the NN interaction. For a quenched weak coupling constant the NMEs are significantly smaller than those for an unquenched g_A mostly due to the factor $(1.00/1.269) = 0.62$ entering in the definition of $M'^{0\nu}$ in Eq. (16). The RQRPA predictions are suppressed when compared to the QRPA ones, especially for ^{76}Ge .

It goes without saying that further progress in the calculation of the $0\nu\beta\beta$ -decay NMEs is required. Thanks to the theoretical efforts made over the last years the disagreement among the different NMEs is now much less severe than it was some years ago. Currently the main issue of interest is that there exists significant disagreement of the ISM results with those of other approaches and the problem of quenching the axial-vector coupling constant. The uncertainty associated with the calculation of the $0\nu\beta\beta$ -decay NMEs can be reduced by suitably chosen nuclear probes. Complementary experimental information from related processes like the $2\nu\beta\beta$ -decay [40], charge-exchange [44] and particle transfer reactions [45] is also very important. The differences among results of various nuclear structure approaches could be understood by performing an anatomy of the $0\nu\beta\beta$ -decay NME [30, 33, 43]. A recent development in the field is encouraging. There is a reason to believe that the uncertainty in the $0\nu\beta\beta$ -decay will be further reduced.

IV. SUMMARY

The possible establishment of CP-violation in the lepton sector is one of the most challenging problems of neutrino and astrophysics. Studies of $0\nu\beta\beta$ -decay driven by Majorana neutrinos can lead to insights into CP violation in this sector. In view of recent measurement of the smallest neutrino mixing angle θ_{13} in the Double Chooz, Daya Bay and RENO experiments we revisited the problem of determining the Majorana CP phases via the additional observation of the $0\nu\beta\beta$ -decay.

Both cases of the normal and inverted hierarchy of neutrino masses were discussed. It was shown that in the case of the NH the determination of one of the Majorana CP phases would be possible only by knowledge of both the absolute value of the effective Majorana mass $|m_{\beta\beta}|$ and of the lightest neutrino mass m_1 . This task cannot be solved by any of the planned or prepared neutrino experiments, only within some particle physics models which allow for a prediction of neutrino masses. Further, it was also found that for some values of m_1 the minimal value of $|m_{\beta\beta}|$ is realized in the case of CP violation.

The case of the IH of neutrino masses offers different possibilities. Future cosmological measurements have the potential to constrain the lightest neutrino mass m_0 to values below 10 meV. The difference of Majorana phases α_{21} depends very weakly on m_0 for these low values and it can be determined by an accurate measurement leading to $|m_{\beta\beta}|$. For this purpose the $0\nu\beta\beta$ -decay NME needs to be evaluated with an uncertainty less than 30%. This is a formidable task, which might be achieved at some point in time due to further developments in the fields of nuclear structure and many-body physics also linked to a further increase of computer power.

ACKNOWLEDGMENTS

This work is supported in part by the Deutsche Forschungsgemeinschaft within the project "Nuclear matrix elements of Neutrino Physics and Cosmology" FA67/40-1 and by the grant of the Ministry of Education and Science of the Russian Federation (contract 12.741.12.0150). F. Š. acknowledges also support by the VEGA Grant agency of the Slovak Republic under the contract No. 1/0639/09.

-
- [1] R. Wendell *et al.* (The Super-Kamiokande Collaboration), Phys. Rev. D **81**, 09200 (2010).
 - [2] B. Aharmim *et al.* (The SNO Collaboration), Phys. Rev. C **81**, 055504 (2010).
 - [3] A. Gando *et al.* (The KamLAND Collaboration), Phys. Rev. D **83**, 052002 (2011).
 - [4] A. Habig *et al.* (The MINOS Collaboration), Mod. Phys. Lett. A **25**, 1219 (2010).
 - [5] B. Pontecorvo, J. Exptl. Theoret. Phys. **33**, 549 (1957). [Sov. Phys. JETP **6**, 429 (1958)]; J. Exptl. Theoret. Phys. **34**, 247 (1958) [Sov. Phys. JETP **7**, 172 (1958)].
 - [6] M. Fukugita, T. Yanagida, Phys. Lett. B **174**, 45 (1986).
 - [7] F.P. An *et al.* (Daya Bay Collaboration), Phys. Rev. Lett. **108**, 171803 (2012).
 - [8] J.K. Ahn *et al.* (RENO Collaboration), Phys. Rev. Lett. **108**, 191802 (2012).
 - [9] K. Abe *et al.* (T2K Collaboration), Phys. Rev. Lett. **107**, 041801 (2011).
 - [10] Y. Abe *et al.* (Double Chooz Collaboration), e-Print: arXiv:1207.6632 [hep-ex].
 - [11] S.M. Bilenky, J. Hosek, S.T. Petcov, Phys. Lett. B **94**, 4 (1980).
 - [12] J. Schechter, J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); M. Doi, et al., Phys. Lett. B **102**, 323 (1981); J. Bernabeu, P. Pascual, Nucl. Phys. B **228**, 21 (1983).
 - [13] P. Langacker, et al., Nucl. Phys. B **282** (1987) 589.
 - [14] V. Barger, et al., Phys. Lett. B **540**, 247 (2002).
 - [15] A. de Gouvea, B. Kayser, R. Mohapatra, Phys. Rev. D **67**, 053004 (2003).
 - [16] S. Pascoli, S.T. Petcov, W. Rodejohann, Phys. Lett. B **549**, 177 (2002).
 - [17] S. Pascoli, S.T. Petcov, T. Schwetz, Nucl. Phys. B **734**, 24 (2006).
 - [18] F.T. Avignone, S.R. Elliott and J. Engel, Rev. Mod. Phys. **80**, 481 (2008).
 - [19] J.D. Vergados, H. Ejiri, and F. Šimkovic, Rep. Prog. Phys. **75**, 096301 (2012).
 - [20] C. Athanassopoulos *et al.*, Phys. Rev. Lett. **75**, 2650 (1995); **77**, 3082 (1996); **81** 1774 (1998). L.B. Auerbach et. al. (LSND Collaboration), Phys. Rev C **64**, 065501 (2001).
 - [21] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration) Phys. Rev. Lett. **102**, 101802 (2009); Z. Djurcic (for the Mini-BooNE Collaboration), e-Print: arXiv:1201.1519 [hep-ex].
 - [22] G. Mention *et al.*, Phys. Rev. D, **83** 073006 (2011).
 - [23] F. Kaether, W. Hampel, G. Heusser, J. Kiko, T. Kirsten, Phys. Lett. B **685**, 47 (2010); J.N. Abdurashitov, V.N. Gavrin, S.V. Girin, V.V. Gorbachev, P.P. Gurkina, T.V. Ibragimova, et al., Phys. Rev. C **80**, 015807 (2009); D. Frekers et al., Phys. Lett. B **706**, 134 (2011).
 - [24] T. Schwetz, M. Tórtola, and J.W.F. Valle, New J. Phys. **10**, 113011 (2008).
 - [25] E.W. Otten and C. Weinheimer, Rep. Prog. Phys. **71**, 086201 (2008).
 - [26] MARE Collaboration, E. Andreotti *et al.*, Nucl. Instrum. Meth. A **572**, 208 (2007); A. Nucciotti, arXiv: 1012.2290 [hep-ex].
 - [27] K.N. Abazajian *et al.*, Astropart. Phys. **35**, 177 (2011).
 - [28] S. A. Thomas, F. B. Abdalla and O. Lahav, Phys. Rev. Lett. **105**, 031301 (2010).
 - [29] V.A. Rodin, A. Faessler, F. Šimkovic, and P. Vogel, Nucl. Phys. A **766**, 107 (2007), Erratum-ibid **793**, 213 (2007).
 - [30] F. Šimkovic, A. Faessler, V.A. Rodin, P. Vogel, and J. Engel, Phys. Rev. C **77**, 045503 (2008).

- [31] F. Šimkovic, A. Faessler, H. Muther, V. Rodin, M. Stauf, Phys. Rev. C **79**, 055501 (2009).
- [32] F. Šimkovic, G. Pantis, J. D. Vergados, and A. Faessler, Phys. Rev. C **60**, 055502 (1999).
- [33] F. Šimkovic, R. Hodák, A. Faessler, and P. Vogel, Phys. Rev. C **83**, 015502 (2009).
- [34] J. Menéndez, A. Poves, E. Caurier, and F. Nowacki, Nucl. Phys. A **818**, 139 (2009).
- [35] M. Horoi and S. Stoica, Phys. Rev. C **81**, 024321 (2010).
- [36] V.A. Rodin, A. Faessler, F. Šimkovic, and P. Vogel, Phys. Rev. C **68**, 044302 (2003);
- [37] P.K. Rath, R. Chandra, K. Chaturvedi, P.K. Raina, J.G. Hirsch, Phys. Rev. C **82**, 064310 (2010).
- [38] T.R. Rodríguez and G. Martínez-Pinedo, Phys. Rev. Lett. **105**, 252503 (2010).
- [39] J. Barea, F. Iachello, Phys. Rev. C **79**, 044301 (2009).
- [40] A.S. Barabash, Phys. Rev. C **81**, 035501 (2010).
- [41] EXO Collaboration (M. Auger (Bern U.) et al.), Phys. Rev. Lett. **109**, 032505 (2012).
- [42] A. Gando *et al.* (The KamLAND-Zen Collaboration), e-Print: arXiv:1205.6130 [hep-ex].
- [43] F. Šimkovic, A. Faessler, and P. Vogel, Phys. Rev. C **79**, 015502 (2009).
- [44] D. Frekers, Prog. Part. Nucl. Phys., **64**, 281 (2010).
- [45] J.P. Schiffer *et al.*, Phys. Rev. Lett. **100**, 112501 (2008).